



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2019

---

## Models of triple covers

Kresch, Andrew ; Tschinkel, Yuri

DOI: <https://doi.org/10.1134/s000143461905016x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-174502>

Journal Article

Accepted Version

Originally published at:

Kresch, Andrew; Tschinkel, Yuri (2019). Models of triple covers. *Mathematical Notes*, 105(5-6):795-797.

DOI: <https://doi.org/10.1134/s000143461905016x>

# MODELS OF TRIPLE COVERS

ANDREW KRESCH AND YURI TSCHINKEL

ABSTRACT. We exhibit, for a degree 3 covering of algebraic varieties, a model where the covering is a finite covering of smooth projective varieties branched over a smooth divisor.

## 1. INTRODUCTION

Let  $k$  be a field of characteristic 0. It is well known that any morphism of projective varieties  $\psi: T \rightarrow S$  over  $k$ , that is generically finite of degree 2, can be put into a commutative diagram

$$\begin{array}{ccc} \tilde{T} & \xrightarrow{\varrho_T} & T \\ \tilde{\psi} \downarrow & & \downarrow \psi \\ \tilde{S} & \xrightarrow{\varrho_S} & S \end{array} \quad (1)$$

with smooth projective varieties  $\tilde{S}$  and  $\tilde{T}$ , such that  $\varrho_S$  is a birational morphism,  $\varrho_T$  is a birational map, and  $\tilde{\psi}$  is a degree 2 finite covering. The branch locus of a degree 2 finite covering of smooth projective varieties is a smooth divisor.

In this note, we establish an analogous theorem for triple covers, over perfect fields of characteristic not equal to 2 or 3. Good models of triple covers are important for the construction of models of fibrations (over a base of arbitrary dimension), when the symmetry group of the geometric generic fiber admits the symmetric group  $\mathfrak{S}_3$  as a quotient, as is the case for fibrations in sextic del Pezzo surfaces [1]–[4].

There is an extensive literature on triple covers of surfaces, e.g., [5]–[9]. In [8], a theorem similar to the one in this note is proved for triple covers of surfaces, by a method related to the classical solution of a cubic equation. Our approach is more geometric, and is based on an analysis of ramification in codimension 1 and 2.

We produce  $\tilde{\psi}: \tilde{T} \rightarrow \tilde{S}$ , ramified over a smooth divisor of  $\tilde{S}$ . By contrast, many of the (non-cyclic) degree 3 coverings of smooth projective varieties that occur naturally have singular branch locus, such as

6-cuspidal sextic with cusps on a conic as branch locus of a general projection of a smooth cubic surface [10]. Such a covering is totally ramified (geometrically, only one pre-image in  $T$ ) only over the cusps; after the procedure described here has been applied there is total ramification over entire components of the branch locus.

As explained in [11, Exa. 3.1], models of degree  $\geq 4$  covers of surfaces as in (1) do not exist in general.

**Acknowledgments:** The second author was partially supported by NSF grant 1601912 and by the Laboratory of Mirror Symmetry NRU HSE, RF Government grant, ag. No. 14.641.31.0001.

## 2. FINITE COVERS AND RAMIFICATION

Let  $k$  be a field and  $S$  a smooth variety over  $k$ , i.e., a separated geometrically integral scheme of finite type over  $k$ . By the normalization of  $S$  in a finite field extension of  $k(S)$  we have a canonical correspondence between finite field extensions of  $k(S)$  and connected normal  $k$ -schemes with finite surjective morphism to  $S$ . The set of points where such a morphism  $T \rightarrow S$  fails to be étale is a closed subset  $Z \subset T$  which is

- equal to  $T$  if and only the associated finite field extension of  $k(S)$  is inseparable;
- is otherwise of pure codimension 1 or empty, by the Zariski-Nagata purity theorem [12, Thm. X.3.1].

Consequently, if there exists a simple normal crossing divisor  $D \subset S$ ,  $D = D_1 \cup \dots \cup D_n$  with  $D_i$  irreducible for all  $i$ , such that  $T \times_S (S \setminus D) \rightarrow S$  is étale, then the branch locus (the image of  $Z$  in  $S$ ) is of the form  $\bigcup_{i \in I} D_i$  for some  $I \subset \{1, \dots, n\}$ .

Once a finite field extension of  $k(S)$  has been specified, when we refer to the branch locus we mean the branch locus of  $T \rightarrow S$ , where  $T$  is the normalization of  $S$  in the given field extension of  $k(S)$ .

## 3. MAIN RESULT

Let  $k$  be a perfect field of characteristic not equal to 2 or 3,  $S$  a smooth projective variety over  $k$ , and  $T$  a projective variety with morphism to  $S$  that is generically finite of degree 3. If the field extension  $k(T)/k(S)$  is cyclic, then an argument just as in the case of double covers yields a commutative diagram (1), where  $\tilde{\psi}: \tilde{T} \rightarrow \tilde{S}$  is a cyclic degree 3 covering of smooth projective varieties branched over a smooth divisor on  $\tilde{S}$ . As in the case of double covers, a form of resolution of singularities for divisors on  $S$  is required; for instance, it is sufficient if the following is available:

(R) embedded resolution of singularities of divisors on  $S$  by iterated blow-up with smooth center.

Of course, this is available (Hironaka) when  $k$  has characteristic zero. When  $k$  has positive characteristic, this is available for  $\dim(S) \leq 3$  (trivial/classical for  $\dim(S) \leq 2$ , due to Abhyankar for  $\dim(S) = 3$ ).

**Theorem 1.** *Let  $k$  be a perfect field of characteristic not equal to 2 or 3,  $S$  a smooth projective variety over  $k$ , and  $\psi: T \rightarrow S$  a morphism of projective varieties that is generically finite of degree 3. We suppose that (R) holds for  $S$ . Then there exists a commutative diagram*

$$\begin{array}{ccc} \tilde{T} & \xrightarrow{\varrho_T} & T \\ \tilde{\psi} \downarrow & & \downarrow \psi \\ \tilde{S} & \xrightarrow{\varrho_S} & S \end{array}$$

with smooth projective varieties  $\tilde{S}$  and  $\tilde{T}$ , such that

- $\varrho_S$  is a birational morphism;
- $\varrho_T$  is a birational map;
- $\tilde{\psi}$  is a degree 3 finite covering, branched over a smooth divisor of  $\tilde{S}$ .

*Proof.* As mentioned, when  $k(T)/k(S)$  is cyclic this is achieved by applying (R) to make the branch locus into a simple normal crossing divisor and repeatedly blowing up components of the intersection of a pair of components of the branch locus to make the branch locus smooth. Essential for the second step is the observation that a component  $Y$  of an intersection  $D_i \cap D_j$  ( $i \neq j$ ) of components of the simple normal crossing branch locus  $D = D_1 \cup \cdots \cup D_n$  (achieved by the first step) may be assigned to one of two types, according to whether the branch locus for  $k(T)/k(S)$  of the blow-up  $Bl_Y S$  has the exceptional divisor as a component. Let  $\ell$ , respectively  $m$  denote the number of components  $Y \subset D_i \cap D_j$  for some  $i \neq j$ , for which the exceptional divisor of  $Bl_Y S$  is, respectively is not, a component of the branch locus. (For simplicity of notation we continue to write  $S$ , even though the first step potentially makes a birational modification.) By blowing up some  $Y$  for which the exceptional divisor is not a component of the branch locus whenever this is possible, and otherwise blowing up any  $Y$ , we obtain for  $(\ell, m) \neq (0, 0)$  a pair  $(\ell', m')$  associated with  $S' := Bl_Y S$  that is smaller than  $(\ell, m)$  in lexicographic order.

In the non-cyclic case we proceed with the same first step, making the branch locus into a simple normal crossing by applying (R). Now the branch locus has components of two kinds: some, say  $D_1 \cup \cdots \cup D_n$ , with

simple ramification and others,  $D'_1 \cup \cdots \cup D'_n$ , with total ramification. We adopt the convention that the  $D_i$  and  $D'_{i'}$  are all irreducible.

The discriminant of  $k(T)/k(S)$  determines a quadratic extension of  $k(S)$  with branch locus  $D_1 \cup \cdots \cup D_n$ . By blowing up intersections of pairs of components, we achieve  $D_i \cap D_j = \emptyset$  for  $i \neq j$ .

We claim that  $D_i \cap D'_{i'} = \emptyset$  for all  $i$  and  $i'$ . This follows from the fact that the pre-image of  $D'_{i'}$  in the discriminant double cover would be irreducible. We argue by contradiction, replacing  $S$  by a strict henselization of the local ring of  $S$  at the generic point of a component of  $D_i \cap D'_{i'}$ . We still have a cubic extension of the residue field of the generic point, still with nontrivial discriminant, and this yields a cyclic cubic extension of the discriminant cover. This must be obtained from a cyclic cubic extension below by base change to the double cover, and we have a contradiction.

We may have  $D'_{i'} \cap D'_{j'} \neq \emptyset$  for some  $i' \neq j'$ , but then as described at the beginning of the proof we may deal with this by blowing up components of intersections  $D'_{i'} \cap D'_{j'}$ . Then we have a smooth branch locus and hence a smooth model of the covering.  $\square$

## REFERENCES

- [1] J.-L. Colliot-Thélène, N. A. Karpenko, and A. S. Merkurjev, Rational surfaces and the canonical dimension of the group  $\mathrm{PGL}_6$ , *Algebra i Analiz*, 19(5):159–178, 2007.
- [2] M. Blunk, Del Pezzo surfaces of degree 6 over an arbitrary field. *J. Algebra*, 323(1):42–58, 2010.
- [3] N. Addington, B. Hassett, Yu. Tschinkel, and A. Várilly-Alvarado, Cubic four-folds fibered in sextic del Pezzo surfaces, [arXiv:1606.05321](#).
- [4] A. Kuznetsov, Derived categories of families of sextic del Pezzo surfaces, [arXiv:1708.00522](#).
- [5] P. du Val, On triple planes having branch curves of order not greater than twelve, *J. London Math. Soc.*, 8(3):199–206, 1933.
- [6] R. Miranda, Triple covers in algebraic geometry, *Amer. J. Math.*, 107(5):1123–1158, 1985.
- [7] H. Tokunaga, Triple coverings of algebraic surfaces according to the Cardano formula, *J. Math. Kyoto Univ.*, 31(2):359–375, 1991.
- [8] S.-L. Tan. Triple covers on smooth algebraic varieties, *Geometry and nonlinear partial differential equations (Hangzhou, 2001)*, *AMS/IP Stud. Adv. Math.*, 29, Amer. Math. Soc., Providence, RI, 2002, 143–164.
- [9] T. Shirane, A note on normal triple covers over  $\mathbf{P}^2$  with branch divisors of degree 6, *Kodai Math. J.*, 37(2):330–340, 2014.
- [10] O. Zariski, On the problem of existence of algebraic functions of two variables possessing a given branch curve, *Amer. J. Math.*, 51(2):305–328, 1929.
- [11] T. Shirane, On 4-fold covers of algebraic surfaces, *Kyushu J. Math.*, 64(2):297–322, 2010.

- [12] A. Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*, Springer-Verlag, Berlin, 1971.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE  
190, CH-8057 ZÜRICH, SWITZERLAND  
*Email address:* `andrew.kresch@math.uzh.ch`

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

SIMONS FOUNDATION, 160 FIFTH AV., NEW YORK, NY 10010, USA  
*Email address:* `tschinkel@cims.nyu.edu`